

DESY 96-233
 hep-ph/9612409
 December 1996

GENTLE/4fan v. 2.0

A Program for the Semi-Analytic Calculation of Predictions for the Process $e^+e^- \rightarrow 4f$

D. Bardin^{a,1,2}, J. Biebel^a, D. Lehner^{b,3},
 A. Leike^{c,4}, A. Olchevski^d, T. Riemann^{a,2}

^a*Deutsches Elektronen-Synchrotron DESY, Institut für Hochenergiephysik
 IfH Zeuthen, Platanenallee 6, D-15738 Zeuthen, Germany*

^b*Fakultät für Physik, Albert-Ludwigs-Universität, Hermann-Herder-Str. 3,
 D-79104 Freiburg, Germany*

^c*Sektion Physik, Ludwig-Maximilians-Universität, Theresienstraße 37,
 D-80333 München, Germany*

^d*Laboratory of Nuclear Problems, JINR, ul. Joliot-Curie 6, RU-141980 Dubna,
 Moscow Region, Russia*

Abstract

We describe version 2.0 of the FORTRAN program **GENTLE/4fan** for the semi-analytic computation of cross-sections and distributions in four-fermion production in e^+e^- annihilation. **GENTLE/4fan** covers all charged current and neutral current four-fermion final states with no identical particles, no electrons, and no electron neutrinos in the final state. Initial state radiation representing the most relevant quantum corrections and anomalous triple gauge boson couplings have been included.

¹ On leave of absence from Bogoliubov Theor. Lab., JINR, ul. Joliot-Curie 6, RU-141980 Dubna, Moscow Region, Russia

² Supported by European Union with grant INTAS-93-744

³ Address after 1 August 1996: INNOSOFT Ltd., Bodenseestraße 4, D-81241 München, Germany

⁴ Supported by European Union with grant CHRX-CT940579

PROGRAM SUMMARY

Title of program: GENTLE/4fan

Version: 2.0 (December 1996)

Catalogue number:

Program obtainable from:

<http://www.ifh.de/theory> or on request by e-mail from the authors

Licensing provisions: non

Computers: all

Operating system: all

Program language: FORTRAN-77

Memory required to execute with typical data: 0.5 Mb

No. of bits per word: 32

No. of lines in distributed program: 8300

Other programs called: none

External files needed: none

Keywords: e^+e^- annihilation, QED, higher order corrections,
four-fermion production, WW , ZZ , $Z\gamma$, $\gamma\gamma$, ZH production

Nature of physical problem:

Description of two-boson production in e^+e^- annihilation at high energies with selected possible background contributions; initial state QED corrections (ISR); selected final state corrections; anomalous triple gauge boson couplings

Method of solution:

Semi-analytic; numerical integration of analytic formulae:

- (i) Born approximation: Numerical integrations over two virtualities s_1, s_2 plus optionally over the boson production angle θ ;
- (ii) QED corrections in the flux function approach (FF): one additional numerical integration over virtuality s' ;
- (iii) QED corrections in the structure function approach (SF): alternatively to (ii), two additional numerical integrations over x_1, x_2

Restrictions on complexity of the problem:

Only selective experimental cuts are possible; the program calculates total cross-sections and the distribution in the boson production angle; background contributions are available only for selected final state configurations; no inclusion of pure weak corrections

Typical running time:

The running time strongly depends upon the options used; one finds e.g.:

Born cross-section on-shell: below one sec., off-shell: few secs.;

LLA QED corrections (FF) on-shell: few secs.; off-shell: few mins.;

LLA QED corrections (SF) on-shell: few secs.; off-shell: several mins.;

complete ISR (FF): on-shell: few mins., off-shell: several hours;

with background: from few secs. (no QED) to several hours (with QED);

with anomalous couplings: from few secs. (no QED) to several mins. (with QED)

Contents

1	Introduction	4
2	Physics Background	4
2.1	Off-shell boson pair production	4
2.2	Background contributions	5
2.3	Initial state QED radiative corrections	7
2.4	Angular distributions	9
2.5	Anomalous couplings	11
3	Description of the Program	11
3.1	Features of GENTLE/4fan	11
3.2	Program input	13
3.3	Routines of GENTLE/4fan	19
3.4	Program output	24
3.5	Use of the program	31
A	Numerical Integration in GENTLE/4fan	33
B	Anomalous Triple-Boson Couplings	34
C	Treatment of Extra Neutral Gauge Bosons	35
D	Options of GENTLE	36
	References	37

LONG WRITE-UP

1 Introduction

With the advent of LEP2, now running above the W pair threshold, e^+e^- collisions have entered their multiple heavy boson production era which will continue with a higher energy linear e^+e^- collider (LC). Among the most interesting processes to be studied in high energy e^+e^- annihilation is boson pair production. As heavy bosons decay, their physics is only accessible through their decay products. Thus, at LEP2 or LC, boson pair production must be accessed through the experimental analysis of four-fermion final states,

$$e^+e^- \rightarrow (W^+W^-, ZZ, Z\gamma, \gamma\gamma, ZH, \dots) \rightarrow f_1\bar{f}_2f_3\bar{f}_4 \quad . \quad (1.1)$$

2 Physics Background

The Standard Model (SM) four-fermion final states produced in e^+e^- annihilation have been classified by several authors [1–4]. Using the semi-analytic approach, we have so far studied the production of four-fermion final states that contain neither identical particles nor electrons or electron neutrinos. These four-fermion final states are printed in **boldface** in tables 1 and 2.

Table 1
Number of Feynman diagrams for ‘CC’ type final states.

	$\bar{d}u$	$\bar{s}c$	$\bar{e}\nu_e$	$\bar{\mu}\nu_\mu$	$\bar{\tau}\nu_\tau$
$d\bar{u}$	43	11	20	10	10
$e\bar{\nu}_e$	20	20	56	18	18
$\mu\bar{\nu}_\mu$	10	10	18	19	9

2.1 Off-shell boson pair production

For semi-analytic tree level calculations (flag **IBORNF=0**), the four-particle phase space is parametrized by the boson scattering solid angle, the boson decay solid products’ angles in the corresponding boson rest frames, and the two final state fermion pair invariant masses s_1 and s_2 [5,6]. After analytic

Table 2

Number of Feynman diagrams for **NC** type final states.

	$\bar{d}d$	$\bar{s}s, \bar{b}b$	$\bar{u}u$	$\bar{e}e$	$\bar{\mu}\mu$	$\bar{\tau}\tau$	$\bar{\nu}_e\nu_e$	$\bar{\nu}_\mu\nu_\mu$	$\bar{\nu}_\tau\nu_\tau$
$\bar{d}d$	4·16	32	4·3	48	24	24	21	10	10
$\bar{u}u$	4·3	4·3	4·16	48	24	24	21	10	10
$\bar{e}e$	48	48	48	4·36	48	48	56	20	20
$\bar{\mu}\mu$	24	24	24	48	4·12	24	19	19	10
$\bar{\nu}_e\nu_e$	21	21	21	56	19	19	4·9	12	12
$\bar{\nu}_\mu\nu_\mu$	10	10	10	20	19	10	12	4·3	6

integration of all angular variables, the total cross-section for double-resonant four-fermion production is given by

$$\sigma^{\text{res}}(s) = \int ds_1 \rho_B(s_1) \int ds_2 \rho_B(s_2) \sigma_0(s; s_1, s_2) \quad (2.1)$$

with the Breit-Wigner densities

$$\rho_B(s_i) = \frac{1}{\pi} \frac{\sqrt{s_i} \Gamma_B}{|s_i - M_B^2 + i s_i \Gamma_B / M_B|^2} \times \text{BR} \xrightarrow{\Gamma_B \rightarrow 0} \delta(s_i - M_B^2) \times \text{BR} \quad (2.2)$$

of the resonant bosons. Explicit formulae for the twofold differential cross-sections in the double-differential approximation $\sigma_0(s; s_1, s_2)$ are found in reference [7] for W pair production, in [3] for Z pair production, and in [8] for associated Higgs production. We emphasize that these twofold differential cross-sections are given by very simple formulae. The Breit-Wigner functions are smoothed by mapping (see appendix A). Flag **IMAP** allows for the **CC** case to use the Breit-Wigner functions directly. In the latter case the program works slower by about a factor of two. We note that in this case the invariant mass distributions are accessible. With flag **INCPRC** one may select ZZ , $\gamma\gamma$, or their common production. For the **CC** case, the on-shell limit may be calculated with flag **IONSHL=0**.

The realization of the various generic functions mentioned here and in the following sections is described in section 3.3.

2.2 Background contributions

Although double-resonant amplitudes are dominant in most relevant cases, single- and non-resonant ‘background’ amplitudes contribute non-negligible

to the cross-sections. Including background Feynman diagrams, one can use the numbers from tables 1 and 2 to introduce a transparent notation [3]:

- CC Processes with final states of type $f_1^u \bar{f}_1^d f_2^u \bar{f}_2^d$ are called *charged current processes*: CC11, CC10, CC09; CC20, CC18.
- NC Processes with final states of type $f_1 \bar{f}_1 f_2 \bar{f}_2$ are called *neutral current processes*: NC32, NC24, NC10, NC06; NC48, NC20, NC21, NC19; NC4·16, NC4·12, NC4·03, NC12; NC4·36, NC4·09.
- mix Processes which may be considered as both CC and NC types are called *mixed processes*: mix43, mix19; mix56. An example for a mixed process is the production of $u\bar{d}d\bar{u} \equiv u\bar{u}d\bar{d}$.

In the NC and mix classes, also Feynman diagrams with Higgs bosons contribute. In addition to the two-boson production classes CC03 [7] (W pair production), NC02 [6] (Z pair production), and NC08 [6] ($ZZ, Z\gamma, \gamma\gamma$ production), GENTLE/4fan is able to perform calculations for CC11, CC10, and CC09 [5] as well as NC32, NC24, NC10, NC06 [9]. This is chosen by flag IPROC. In the above NC cases, also Higgs diagrams are included [8].

Complete tree level four-fermion production cross-sections are given by the generic formula

$$\sigma^{\text{Born}}(s) = \int ds_1 \int ds_2 \frac{\sqrt{\lambda}}{\pi s^2} \sum_k \frac{d^2 \sigma_k(s; s_1, s_2)}{ds_1 ds_2} \quad (2.3)$$

with $\lambda \equiv \lambda(s; s_1, s_2)$, $\lambda(a, b, c) = a^2 + b^2 + c^2 - 2ab - 2ac - 2bc$ and

$$\frac{d^2 \sigma_k}{ds_1 ds_2} = \mathcal{C}_k(s; s_1, s_2) \mathcal{G}_k(s; s_1, s_2) \quad (2.4)$$

The coefficient functions $\mathcal{C}_k$ represent coupling constants and boson propagators, while $\mathcal{G}_k$ is a kinematical function obtained after analytical performance of angular integration. The index k labels cross-section contributions with both different coupling structure and different Feynman topology. Noticing that often different twofold differential cross-sections $d^2 \sigma_k / ds_1 ds_2$ are related by symmetries in their arguments, one realizes that equation (2.3) enables efficient coding of the cross-section. For CC processes, background contributions are switched off with flag IBCKGR=0. With flag ICHNNL one may choose the final state topology in the CC case and with IFERM1 and IFERM2 in the NC case.

2.3 Initial state QED radiative corrections

In e^+e^- annihilation, the most relevant radiative corrections are initial state QED corrections (ISR). The phase space is now parameterized by the center-of-mass photon scattering solid angle, by the boson scattering solid angle in the two-boson (or, equivalently, four-fermion) rest frame, by the boson decay products' solid angles in the corresponding boson rest frames, by the final state fermion pair invariant masses, and by the four-fermion (or reduced) invariant mass s' [6]. After analytic integration over all angular degrees of freedom, the $\mathcal{O}(\alpha)$ ISR-corrected total four-fermion production cross-section with soft photon exponentiation is determined in the flux function approach (flag ICONVL=0) by

$$\sigma^{\text{ISR}}(s) = \int ds_1 \int ds_2 \int ds' \sum_k \frac{d^3\sigma_k(s, s'; s_1, s_2)}{ds_1 ds_2 ds'} , \quad (2.5)$$

with

$$\frac{d^3\sigma_k(s, s'; s_1, s_2)}{ds_1 ds_2 ds'} = \mathcal{C}_k \left[\beta_e v^{\beta_e-1} \mathcal{S}_k + \mathcal{H}_k \right] , \quad (2.6)$$

where $\beta_e = 2(\alpha/\pi)(L-1)$, $L \equiv \ln(s/m_e^2)$. The soft+virtual and hard corrections $\mathcal{S}_k$ and $\mathcal{H}_k$ separate into a universal, factorizing, process-independent and a non-universal, non-factorizing, process-dependent part. The latter is included with flag ITNONU=1. They are given by [6,10]

$$\begin{aligned} \mathcal{S}_k &= \left[1 + \bar{S}(s) \right] \sigma_{k,0}(s'; s_1, s_2) + \sigma_{\hat{S},k}(s'; s_1, s_2) , \\ \mathcal{H}_k &= \underbrace{\bar{H}(s, s') \sigma_{k,0}(s'; s_1, s_2)}_{\text{Universal Part}} + \underbrace{\sigma_{\hat{H},k}(s, s'; s_1, s_2)}_{\text{Non-universal Part}} \end{aligned} \quad (2.7)$$

with $\sigma_{k,0}(s'; s_1, s_2) \equiv \left[\sqrt{\lambda}/(\pi s'^2) \right] \mathcal{G}_k(s'; s_1, s_2)$. The universal $\mathcal{O}(\alpha)$ soft+virtual and hard radiators $\bar{S}$ and $\bar{H}$ read

$$\bar{S}(s) = \frac{\alpha}{\pi} \left(\frac{\pi^2}{3} - \frac{1}{2} \right) + \frac{3}{4} \beta_e , \quad (2.8)$$

$$\bar{H}(s, s') = -\frac{1}{2} \left(1 + \frac{s'}{s} \right) \beta_e . \quad (2.9)$$

Flag IZERO influences $\bar{S}(s)$. If the index k labels pure s-channel e^+e^- annihilation, the non-universal contribution vanishes. Non-universal ISR contributions are analytically very complex, but, in contrast to the universal corrections, not

logarithmically enhanced.

For the CC03 process, a significant QED radiative correction is the so-called Coulomb singularity [11,12,14] originating from the exchange of photons between the two resonant W bosons. It effects in a factor $(1 + \mathcal{F}_{Coulomb})$ to be multiplied to the twofold differential cross-section of equation (2.4) or to the threefold differential cross-section of equation (2.6). Details are defined with flag **ICOLMB**.

The leading logarithmic ISR contributions (essentially what we called the universal ISR contributions) may also be determined using the so-called structure function approach [15] (flag **ICONVL=1**). The basic formula, adapted for our case, is

$$\frac{d^2\sigma_{e^+e^- \rightarrow 4f n\gamma}}{ds_1 ds_2} = \int_{x_+^{\min}}^1 dx_+ \int_{x_-^{\min}}^1 dx_- D(x_+)D(x_-) \sigma_0(s'; s_1, s_2) \quad (2.10)$$

with the tree level twofold differential cross-section σ_0 evaluated at $s' = sx_+x_-$. The variables x_+ and x_- represent the momentum fractions of the e^+ and e^- after the radiation of photons. Their minimum values are given by

$$\begin{aligned} x_+^{\min} &= \frac{(\sqrt{s_1} + \sqrt{s_2})^2}{s}, \\ x_-^{\min} &= \frac{(\sqrt{s_1} + \sqrt{s_2})^2}{x_+ s}. \end{aligned} \quad (2.11)$$

An exhaustive discussion of the application of the structure function approach to $4f$ production may be found in [1,15] and in references quoted therein. Our implementation of the structure functions $D(x)$ follows equations (4.18) to (4.21) of reference [5]. Note that, to improve the calculation, σ_0 can be replaced by a “dressed” tree level cross-section which includes the running QED coupling and/or the Coulomb factor. Since the soft-photon pole is proportional to $(L - 1)$ rather than to the leading logarithm L alone there are the so-called **BETA** and **ETA** alternatives for the implementation of the structure functions [15]. They are selected by flag **IZETTA**.

For both descriptions of QED corrections, the details of higher order terms are defined by flag **IQEDHS**.

In addition to the total cross-section, **GENTLE/4fan** can compute the angular distribution, bin-integrated cross-sections, and moments for several other physically relevant quantities. These quantities are selected with the variables **IDCS** and **IREGIM**, the latter being chosen with flags **IRMAX**, **IRSTP**:

- (1) the radiative final state four-fermion invariant mass loss $X = (s-s')/(2\sqrt{s})$ for the **CC11** family in the flux function as well as the structure function approach;
- (2) the true radiative energy loss $X = \sqrt{s}/2 [(1-x_+) + (1-x_-)]$ for the **CC11** family in the structure function approach;
- (3) the W mass shift $X = [(\sqrt{s_1} + \sqrt{s_2})/2 - M_W]$ for the **CC11** family in the flux function as well as the structure function approach;
- (4) moments X related to the angular distribution are treated in section 2.4.

The running of loop variable **IMOMN**= n is defined by flags **IMMIN**, **IMMAX**. It is used for the selection of the moment of n^{th} degree $M_{(X,n)}$ of a quantity X (calculated with QED corrections):

$$M_{(X,n)} = \frac{\int X^n \sigma_X}{\int \sigma_X} \quad (2.12)$$

where X is selected by **IREGIM** and the integration is over the phase space variables upon which X depends and σ_X is the cross-section differential in exactly those phase space variables. Note that $M_{(X,1)}$ is an average of the quantity X .

Finally, we mention that an inclusive implementation of QCD corrections is provided. Firstly, QCD corrections to the total W width have been implemented by a factor $(1 + 2\alpha_s/3\pi)$ to be multiplied to Γ_W . Secondly, the final state radiative QCD corrections to the formation of a quark-anti-quark pair are built in by multiplying the relevant branching ratio with $(1 + \alpha_s/\pi)$.

2.4 Angular distributions

Because of the importance of the **CC03** process, **GENTLE/4fan** is also able to compute W production angular distributions, bin-integrated cross-sections and moments for this process. A selection is possible by the flags **IDCS** and **IREGIM**. In the first and last of the three mentioned cases, the last analytical integration is not carried out. Instead, in Born approximation the analytic threefold differential cross-section

$$\frac{d^3\sigma_{\text{CC03}}}{ds_1 ds_2 d\cos\theta} = \frac{\sqrt{\lambda}}{\pi s^2} \sum_{k=1}^3 C_k^{\text{CC03}} \mathcal{G}_k^{\text{CC03}}(s, s_2, s_1; \cos\theta) \quad (2.13)$$

is used. The angular kinematic functions $\mathcal{G}_k^{\text{CC03}}(s, s_2, s_1; \cos\theta)$ were published in [5]. The cross-section $d\sigma_{\text{CC03}}/d\cos\theta$ is obtained upon integration over s_1 and s_2 . In **GENTLE/4fan**, as a measure for the **CC03** angular distribution, the first four moments of the differential cross-section (2.13) may be calculated as

its convolution with Chebyshev polynomials $T_n(\cos\theta)$. The Born moment of n^{th} degree is given by

$$M_{(\cos\theta,n)} = \int ds_1 ds_2 d\cos\theta T_n(\cos\theta) \frac{d^3\sigma_{cc03}}{ds_1 ds_2 d\cos\theta} . \quad (2.14)$$

The zeroth moment is just the total cross-section.

Initial state QED corrections to the angular distributions and moments have to be calculated in the structure function approach. With photon emission, the W production angle in the laboratory system, θ_{lab} , and in the center-of-mass system of the four final state fermions, θ , are different. The Lorentz boost relating them depends on both x_+ and x_- . The QED corrected angular moments with ISR are given by

$$M_{(\cos\theta_{\text{lab}},n)} = \int ds_1 \int ds_2 \int dx_+ \int dx_- \int d\cos\theta \\ \times D(x_+)D(x_-) T_n[\cos\theta_{\text{lab}}(\cos\theta)] \frac{d^3\sigma_{cc03}}{ds_1 ds_2 d\cos\theta} . \quad (2.15)$$

With flags IDCS, IBIN, IBINNU, the user may select the calculation of the angular distribution:

$$\frac{d\sigma_{cc03}}{d\cos\theta_{\text{lab}}} = \int ds_1 \int ds_2 \int dx_+ \int dx_- \\ \times D(x_+)D(x_-) \sum_i \left| \frac{\partial \cos\theta_i}{\partial \cos\theta_{\text{lab}}} \right| \frac{d^3\sigma_{cc03}}{ds_1 ds_2 d\cos\theta_i} \quad (2.16)$$

with the threefold differential tree level cross-section $d^3\sigma_{cc03}/(ds_1 ds_2 d\cos\theta_i)$ depending on $s' = sx_+x_-$, s_1 , s_2 , and $\cos\theta_i(s, s_1, s_2, x_+, x_-, \cos\theta_{\text{lab}})$. Index i indicates that there may be zero, one, or two solutions for the Lorentz boost.

In analogy to (2.15), the bin-integrated cross-section formulae may be written with the boosted angular argument under the integral, thus allowing for relatively simple analytic angular integrations (and avoiding the appearance of the Jacobean) if, of course, the integration limits set in the laboratory frame are transformed into angular limits in the boosted frame. In this respect, the approach to the bin-integrated cross-section differs from that for the angular distribution where we integrate over the laboratory angle (and have to take into account the Jacobean). A choice can be made with flag IBIN.

2.5 Anomalous couplings

Angular distributions as defined in the above section 2.4 are especially useful in the context of anomalous triple boson couplings [16]. In **GENTLE/4fan**, only those anomalous $\gamma W^+ W^-$ and $Z W^+ W^-$ couplings are implemented which obey Lorentz invariance, conserve **CP**, and do not modify the electromagnetic interaction. These constraints allow a restricted set of anomalous contributions to the triple gauge boson vertex. These couplings are denoted by $g_3^a(V)$ with $V \in \{\gamma, Z\}$. The $a = s, x, y, z$ denotes the coupling type including the **SM** triple-boson coupling $g_3^s(V)$. The couplings are derived from the anomalous couplings defined in equations (B.3) to (B.8). The threefold differential **CC03** cross-section is then given by [17]

$$\frac{d^3 \sigma_{\text{CC03}}^{\text{ano}}}{ds_1 ds_2 d \cos \theta_{\text{lab}}} = \frac{\sqrt{\lambda}}{\pi s^2} \left(\mathcal{C}^t \mathcal{G}^t + \sum_k \mathcal{C}_k^{st} \mathcal{G}_k^{st} + \sum_{ij} \mathcal{C}_{ij}^s \mathcal{G}_{ij}^s \right) , \quad (2.17)$$

where i, j , and k run over s, x, y, z . The summation over s-channel photon and Z amplitudes is contained in the coefficient functions $\mathcal{C}$. Compared to the **SM** where only three kinematical functions describe the **CC03** process, the inclusion of anomalous couplings adds considerable complexity to the calculation. The user may select a treatment of anomalous couplings with flag **IANO**. The potential influence of an extra heavy neutral boson Z' is explained in appendix C.

3 Description of the Program

GENTLE/4fan is a **FORTRAN77** program for the computation of total and differential cross-sections for four-fermion production in $e^+ e^-$ annihilation with inclusion of the relevant initial state QED corrections. It originated from older versions [18].

3.1 Features of **GENTLE/4fan**

The cross-sections, angular distributions, and moments presented in equations (2.1), (2.3), (2.5), (2.10), (2.12), (2.14), (2.15), (2.16), and (2.17) are computed in **GENTLE/4fan** by numerically integrating analytical formulae for expressions such as equation (2.6). The analytical input formulae are found in [3,5,7–9,6,10–12,17] and in references therein. Numerical integrations are performed by a fast and stable self-adaptive Simpson algorithm realized in the

subroutines `SIMPS` and `FDSIMP`⁵.

For transparency – and also for historical reasons – version 2.0 of `GENTLE/4fan` is built from three pieces of code. The corresponding ramification is controlled by the flag `IPROC`. The first part contains the charged current (`CC`) reactions including $\mathcal{O}(\alpha)$ initial state QED corrections and anomalous couplings. The second part is the `4fan` code with `NC` reactions. The third part, steered by the subroutine `NCGENT` contains the code for the `NC02` and `NC08` processes with complete $\mathcal{O}(\alpha)$ initial state QED corrections. Subsequently we will call the first part the `CC` part, the second one will be named the `4fan` branch, and the third part will be called `NCqed` part. The first and second part together are referred to as the `CC/NC` branch.

At present, the `GENTLE/4fan` may treat the following four-fermion final states:

- (1) `CC03` (with complete ISR, angular distributions, and anomalous couplings) [6,10,17]
- (2) `NC02`, `NC08` (with complete ISR) [6]
- (3) `CC09`, `CC10`, `CC11` [5]
- (4) `NC06`, `NC10`, `NC24`, `NC32` [9]
- (5) `NC` of (4) + Higgs [8]

Initial state radiation is implemented. For total cross-sections *universal* initial state radiative corrections as described in equations (2.5) to (2.7) are computed. In addition, *non-universal* ISR is available for the `CC03`, `NC02`, and `NC08` processes. For the `CC11` family, leading logarithmic initial state QED corrections can also be calculated in the structure function approach as indicated in eq. (2.15). Further, the Coulomb correction to `CC03` and inclusive QCD corrections (to both the denominators of the Breit-Wigner functions and final states with quarks) have been implemented. The user can choose between the different prescriptions of references [11,12] for the Coulomb correction via the flag `ICOLMB`.

Cuts may be applied on the invariant masses s_1 and s_2 . In the structure function approach, cuts on the electron and positron momentum fractions after initial state radiation may be imposed. In the `NCqed` branch of the code, a minimum value for the fraction s'/s may be set as well as for s_1/s and s_2/s . A cut on the `CC03` W production angle is not directly supported by the code but may be imposed using the option for bin-integration.

In general, final state masses are neglected in `GENTLE/4fan`, i.e. the program uses the small mass approximation. Where needed, however, masses are retained in the phase space factors. Masses of heavy particles coupled to the

⁵ The FORTRAN code `FDSIMP` by Yu. Sedych is an extension of the `SIMPS` code by I.N. Silin.

Higgs boson are taken into account where appropriate. In the **NCqed** branch of the program, masses may be chosen by altering the values of **AM1** or **AM2** in the initialization routine **NCIN00**.

3.2 Program input

The common units throughout the code are GeV and GeV/ c^2 . The physical input parameters of **GENTLE/4fan** are defined in subroutine **WUFLAG**:

ALFAI	=	$\alpha(0)$	=	137.0359895, the inverse running fine structure constant at zero momentum transfer
ALFAS	=	$\alpha_s(\text{low } q^2)$	=	0.2
ALPHS	=	$\alpha_s(2M_W)$	=	0.12
ALPW	=	$\alpha(2M_W)$	=	1/128.07, the running fine structure constant at $2M_W$
AME	=	m_e	=	$0.51099906 \times 10^{-3}$ GeV, the electron mass
AMZ	=	M_Z	=	91.1888 GeV, the Z mass
AMW	=	M_W	=	80.230 GeV, the W mass
GAMZ	=	Γ_Z	=	2.4974 GeV, the Z width
GFER	=	G_μ	=	1.16639×10^{-5} GeV $^{-2}$, the Fermi constant ⁶

For the fermion-boson couplings, the value $\alpha(2M_W)$ is used, while for the ISR corrections, the Thompson limit $\alpha(0)$ of the fine structure constant is applied. Depending on the flag value **IINPT**, the weak mixing angle is computed as

$$\text{SIN2W} = \sin^2 \theta_W = \frac{\pi \alpha(2M_W)}{\sqrt{2} M_W^2 G_\mu} \quad \text{for } \text{IINPT} = 0 \quad (3.1)$$

$$\text{SIN2W} = \sin^2 \theta_W = 1 - M_W^2/M_Z^2 \quad \text{for } \text{IINPT} = 1 \quad (3.2)$$

$$\text{SIN2W} = \sin^2 \theta_W = 0.22591 \text{ (numerical user input) for } \text{IINPT} = 2 \quad (3.3)$$

For charged current reactions, the following derived quantities are used in

⁶ $\alpha_s(2M_W)$ is used for the QCD corrections to the W width and to the final state quark-anti-quark pairs in **CC** processes. The low momentum transfer value $\alpha_s(|q| \ll 2M_W)$ is used for the **NC32** process [19] in the calculation of gluon exchange diagrams.

GENTLE/4fan:

$$\text{GAMW} = \Gamma_W = \frac{9}{6\sqrt{2}\pi} G_\mu M_W^3 \left(1 + \frac{2\alpha_s(2M_W)}{3\pi} \right) \text{ for } \text{IGAMW} = 0 \quad (3.4)$$

$$\text{GAMW} = \Gamma_W = 2.085 \text{ (numerical user input)} \quad \text{for } \text{IGAMW} = 1 \quad (3.5)$$

as well as

$$\text{GAE} = -\frac{e}{4s_W c_W} = -\frac{\sqrt{4\pi\alpha(2M_W)}}{4s_W c_W}$$

$$\text{GVE} = \text{GAE}(1 - 4s_W)$$

$$\text{GWF} = \frac{g}{2\sqrt{2}} = -\text{GAE}\sqrt{2}c_W$$

$$\text{GWWG} = \sqrt{4\pi\alpha(2M_W)}$$

$$\text{GWWZ} = \text{GWWG} \frac{c_W}{s_W} .$$

GVE and **GAE** are the electron vector and axial vector couplings, **GWF** is the fermion- W coupling, and **GWWG** and **GWWZ** are the triple gauge boson couplings for the photon and the Z boson in the Standard Model, respectively. All other couplings are uniquely determined by the above. Although, for historical reasons, some of the variable names are slightly different, the same derived quantities are used for neutral current reactions. For the neutral current processes, the final state fermion types and their masses are part of the input. For the **NC08** family of processes, this input is set in the subroutine **NCIN00**. The fermion types for the **NC32** family are set by the flags **IFERM1** and **IFERM2**, while the corresponding masses are loaded from a **DATA** statement in the subroutine **bbmmin**⁷. In the subroutine **WUFLAG**, the Higgs mass **AMHIG** and the Higgs width **GAMHIG** as well as the flags which steer the program flow are initialized. The Higgs contribution may be switched on or off by setting flag **IHIGGS**.

The flags and, for the **CC** branch, the physical input may be overridden by

⁷ Note that the fermion masses are stored in the variables **AM1** and **AM2** for the **NC08** as well as the **NC32** families. This is, however, unproblematic, because the routines for each family are completely separate and are never executed together in a single run.

calls to the subroutine WUFLAG; see also section 3.5. *We strongly recommend the user to carefully control that only options are active which are definitely declared to be compatible with each other.*

In WUFLAG, the following flags are initialized:

IANO

- IANO: to set the anomalous couplings defined in equations (B.3) to (B.8) of appendix B
- IANO=0: all couplings at their SM value
- IANO=±1: $XG = x_\gamma = \pm 0.5$, all other couplings at their SM values
- IANO=±2: $YG = y_\gamma = \pm 0.5$, all other couplings at their SM values
- IANO=±3: $XZ = x_Z = \pm 0.5$, all other couplings at their SM values
- IANO=±4: $YZ = y_Z = \pm 0.5$, all other couplings at their SM values
- IANO=±5: $ZZ = z_Z = \pm 0.1$, all other couplings at their SM values
- IANO=±6: $DZ = \delta_Z = \pm 0.5$, all other couplings at their SM values
- IANO=±7: “HISZ scenario” [20] with
 $x_\gamma = \pm 0.1$,
 $x_Z = x_\gamma (\cos^2 \theta_W - \sin^2 \theta_W) / (2 \sin \theta_W \cos \theta_W)$,
 $\delta_Z = x_\gamma / (2 \sin \theta_W \cos \theta_W)$,
and $y_\gamma = y_Z = z_Z = 0$
- IANO=8: user-set values of x_γ , y_γ , x_Z , y_Z , z_Z , δ_γ , and δ_Z ; for extra Z' couplings, see appendix C

IBCKGR

- IBCKGR: only active for IPROC=1
- IBCKGR=0: calculations for the double resonant CC03 process only, no CC11 background
- IBCKGR=1: inclusion of CC11 background in accordance with ICHNNL not available for IDCS=1

IBIN

- IBIN: only active for IDCS=1
- IBIN=0: calculation of $d\sigma/d\cos\theta$ for IBINNU fixed values of $\cos\theta$
- IBIN=1: calculation of $d\sigma/d\cos\theta$ integrated over each of the IBINNU $\cos\theta$ -bins

IBINNU

- IBINNU: only active for IDCS=1
- IBINNU=n: number of points or bins for the differential cross-section calculation

IBORNF

- IBORNF=0: Born approximation, no radiative corrections
- IBORNF=1: initial state QED corrections are included

ICHNNL

- ICHNNL: only active for IPROC=1 and IBCKGR=1
- ICHNNL=0: no background taken into account, calculation of inclusive CC03 quantities
- ICHNNL=1: calculation of CC09 quantities for a leptonic final state
- ICHNNL=2: calculation of CC10 quantities for a semi-leptonic final state with positively charged lepton
- ICHNNL=3: calculation of CC10 quantities for a semi-leptonic final state with negatively charged lepton
- ICHNNL=4: calculation of CC11 quantities for a hadronic final state
- ICHNNL=5: calculation of inclusive quantities for the CC11 family

ICOLMB

- ICOLMB: determines, how the Coulomb correction is included in the differential cross-sections of equations (2.4) or (2.6) only relevant for the CC03 process
- ICOLMB=0: Coulomb correction not included
- ICOLMB=1: Coulomb correction as in equation (5) of reference [11], but without the phase factor
- ICOLMB=2: Coulomb correction as in equation (5) of reference [11], but with $\Delta \equiv (s_1 - s_2)/s' = 0$
- ICOLMB=3: Coulomb correction as in equation (5) of reference [11]
- ICOLMB=4: Coulomb correction as in reference [14]
- ICOLMB=5: Coulomb correction as in reference [12]

ICONVL

- ICONVL: only active for IPROC=1 and IPROC=2
- ICONVL=0: flux function convolution as given in equation (2.5)
- ICONVL=1: structure function convolution as indicated in eq. (2.10) attention: extremely CPU time consuming

IDCS

- IDCS: only active for IPROC=1 and IBCKGR=0
- IDCS=0: calculation of total cross-sections and moments
- IDCS=1: calculation of the differential cross-section $d\sigma/d\cos\theta$ with anomalous couplings as determined by the flag IANO if used in conjunction with ISR corrections, the structure function approach ICONVL=1 must be used needs IBCKGR=0

IFERM1, IFERM2

IFERM1: only active for IPROC=2 IFERM1 is an integer specifying the first final state fermion pair according to the PDG particle numbering scheme [13]
IFERM2: like IFERM1, but for the second final state fermion pair remember that it is the second final state fermion pair that couples to the Higgs boson, if a calculation with the Higgs is requested

IGAMW

IGAMW=0: computation of the W width according to (3.4)
IGAMW=1: numerical user input

IGAMZS

IGAMZS=0: constant Z width for Z propagators
IGAMZS=1: s -dependent Z width for Z propagators

IHIGGS

IHIGGS: only active for IPROC=2
IHIGGS=0: Higgs not included
IHIGGS=1: Higgs included

IINPT

IINPT=0: weak mixing angle defined by equation (3.1)
IINPT=1: weak mixing angle defined by equation (3.3)
IINPT=2: numerical user input, only active for IPROC=1

IIQCD

IIQCD: only active for IPROC=1
IIQCD=0: no QCD corrections, i.e. $\alpha_s(2M_W) \equiv 0$
IIQCD=1: inclusive QCD corrections to the W width and final state QCD corrections to the final state quark pair are taken into account with $\alpha_s(2M_W) = 0.12$

IMOMN

IMOMN: only active for IPROC=1 and IDCS=0
IMOMN= n : the n^{th} moment of the physical quantity determined by the current IREGIM is computed

The moments to be computed are determined by the values of IMMIN and IMMAX which control the scope of the FORTRAN loop DO IMOMN=IMMIN, IMMAX .

INCPRC

INCPRC: only active for IPROC=3
INCPRC=0: NC02 cross-section calculation, i.e. Z pair production only
INCPRC=1: photon pair production only
INCPRC=2: NC08 cross-section calculation

IONSHL

IONSHL: only active for IPROC=1
IONSHL=0: on-shell limit for W pair production
IONSHL=1: off-shell W bosons

IPROC

IPROC=1: CC11 family of processes
IPROC=2: NC32 family of processes
IPROC=3: NC08 family of processes with complete ISR

IQEDHS

IQEDHS: determines the soft+virtual and hard radiators $\bar{S}$ and $\bar{H}$ to be used for universal corrections to total cross-sections only active, if IBORNF=1 and ICONVL=0
IQEDHS=0: $\mathcal{O}(\alpha)$ exponentiated universal ISR corrections as given in equation (2.6) with the radiators $\bar{S}$ and $\bar{H}$ from equations (2.8) and (2.9)
IQEDHS=1: as IQEDHS=0, but terms of $\mathcal{O}[(\alpha L)^2]$ added to $\bar{S}$ and $\bar{H}$
IQEDHS=2: as IQEDHS=1, but terms of $\mathcal{O}(\alpha^2 L)$ added to $\bar{S}$ and $\bar{H}$
IQEDHS=3: as IQEDHS=2, but terms of $\mathcal{O}(\alpha^2)$ added to $\bar{S}$ and $\bar{H}$

IREGIM

IREGIM: only active for IPROC=1 and IDCS=0
determines the physical quantity to be computed in addition to the total cross-section (see sections 2.3 and 2.4)
IREGIM=0: computation of the total cross-section, used as normalization for next cases
IREGIM=1: computation of moments of the radiative invariant mass loss
IREGIM=2: computation of moments of the radiative energy loss for the structure function approach, i.e. for ICONVL=1 only
IREGIM=3: computation of moments of the W mass shift for the CC03 process
IREGIM=4: computation of W production angular moments for the CC03 process with Chebyshev polynomials (compare section 2.4)

The physical quantities to be computed are determined by the integer values of the variables IRLMAX and IRLSTP which control the scope of IREGIM via the FORTRAN statement DO IREGIM=0,IRLMAX,IRLSTP.

ITNONU

- ITNONU: only active for IBORNF=1
ITNONU=0: only universal hard ISR corrections are taken into account (compare equations (2.7) and (2.9))
ITNONU=1: complete universal and non-universal hard ISR corrections are taken into account; only in effect for CC03 and for the NC08 family, IBCKGR=IDCS = 0
attention: very CPU time consuming

IZERO

- IZERO: only for the flux function convolution, i.e. for ICONVL=0
IZERO=0: the constant term $(\alpha/\pi)(\pi^2/3 - 1/2)$ in the $\bar{S}$ is neglected (compare equation (2.8))
IZERO=1: the soft+virtual $\bar{S}$ radiator is given by equation (2.8)

IZETTA

- IZETTA: only for the structure function convolution, i.e. for ICONVL=1
compare equation (4.21) of reference [5]
IZETTA=0: ETA choice, i.e. in the hard photon radiation part of the structure function $(2\alpha/\pi)L$ is used, whereas $(2\alpha/\pi)(L - 1)$ is entered into the soft part and the exponentiation
IZETTA=1: BETA choice, i.e. $(2\alpha/\pi)(L - 1)$ is used for the whole structure function

Finally, we mention that the relative and absolute precision requirements for the numerical integrations are set at different places in **GENTLE/4fan**. Sensible default values for the precision requirements are chosen for the different branches of the program. In addition, we collect here the physical input parameters which can be chosen by calls to subroutine **WUFLAG**:

ALPHS, ALPW, AME, AMFER2, AMHIG, AMW, AMZ, DZ, GAMHIG, GAMW, GAMZ, GFER, SINW2, XG, XZ, YG, YZ, ZZ.

3.3 Routines of GENTLE/4fan

In this subsection we give a brief description of the subroutines and functions in the code. **GENTLE/4fan** uses **REAL*8** and **COMPLEX*16** variables.

- **FUNCTION alamk** – computes the kinematic λ -function $\lambda(x, y, z) = x^2 + y^2 + z^2 - 2xy - 2xz - 2yz$
- **FUNCTION ALRUN** – may be used in subroutine **prop** to compute photon propagators with running fine structure constant α

- SUBROUTINE `bbmmin` – constitutes the central routine for the `4fan` part of the code, i.e. for computations for the NC32 family; performs the initialization of physical and coupling constants for the `4fan` part
- FUNCTION `c222` (`c322`, `c322m`, `c422`, `c233`) – computes the coupling function $\mathcal{C}_{222}$ ($\mathcal{C}_{322}$, $\mathcal{C}_{322}^a$, $\mathcal{C}_{422}$, $\mathcal{C}_{233}$) introduced in references [8,9]⁸
- FUNCTION `cg422` (`cg233`) – computes the coupling function similar to $\mathcal{C}_{422}$ ($\mathcal{C}_{233}$), but with a gluon exchange replacing the electroweak boson exchange in the final state (compare figure 1e of reference [19])
- SUBROUTINE `check` – checks and prints the input to the `4fan` part; produces the integer error flag `ierr`; prints messages if input errors are encountered
- FUNCTION `COULOM` – computes, according to the flag value of `ICOLMB`, the term `COULOM` for the Coulomb correction factor $(1+\text{COULOM})$
- FUNCTIONS `DDILOG`, `DLI2` – provide the dilogarithm function $\text{Li}_2(x)$ for real argument; if $x \geq 1$, x is understood to be the real part of the argument with infinitesimal imaginary part, and the real part of the function value is returned
- SUBROUTINE `DELTPR` – computes, for the flux function approach, the three-fold (with respect to s_1 , s_2 , and s') differential, non-universal contributions to the CC03 cross-section
- SUBROUTINE `DELVNU` – computes the non-universal soft+virtual contribution to the NCqed cross-section
- FUNCTION `DLI3` – provides the trilogarithm function $\text{Li}_3(x)$ for $x \leq 1$
- FUNCTION `DS12` – provides the Nielsen function $S_{1,2}(x)$ for $x \leq 1$
- SUBROUTINE `dsdshsz` – provides the twofold differential cross-section $d^2\sigma/(ds_1 ds_2)$ for the NC32 family as given in references [8,9]
- FUNCTION `DXS2` – provides, for the CC/NC branch of the code and according to the flags `IONSHL`, `IPROC`, `IBORNF`, `IBCKGR`, and `ICHNNL`, twofold differential expressions (with respect to s_1 and s_2) for the cross-section or for moments by calling `XSECB` and `XSECBG` or `XSEC2E`
- FUNCTION `FACINC` – is the inverse function of `FACNC`
- FUNCTION `FACINV` – represents the inverse function of `FACT`
- FUNCTION `FACNC` – is the mapping function for the s' integrations used in calls to `FDSIMT`
- FUNCTION `FACT` – is the mapping function for the s' , x_+ , and x_- integrations used in calls to `FDSIMP` and `FDSIMS`
- FUNCTION `FCONC` – provides the integrand of the s' integration in `XSNC2E`
- FUNCTION `FCONV1` – provides, according to the flag `ICONVL`, threefold differential (with respect to s_1 , s_2 , and s' or x_+) expressions for
 - (1) the ISR-corrected cross-section, the radiative invariant mass loss, and the W mass shift using the flux function approach
 - (2) total and differential cross-sections as well as moments in the structure function approach by calling `FCONV2`

⁸ Beware: The normalization differs from the references in some cases.

- FUNCTION FCONV2 – provides fourfold differential (with respect to s_1 , s_2 , x_+ , and x_-) expressions for all quantities available in the structure function approach
- SUBROUTINE FDSIMP – is similar to SIMPS, but performs the integration with the help a mapping function which is called in the argument list
- SUBROUTINES FEYNINT1, FEYNINT2, FEYNINT3 – compute, from the analytical formulae of reference [6], the values of several complicated integrals needed for the non-universal hard ISR correction
- FUNCTION FINE12 (FI12NC) – provides the differential cross-section $d\sigma/dx_1$, $x_i = s_i/s$ for the CC/NC (NCqed) branch of the code
- FUNCTION FINE34 (FI34NC) – provides the differential cross-section $d\sigma/(dx_1 dx_2)$ for the CC/NC (NCqed) branch of the code by calling the function DXS2 (XSBRNC and XSNC2E) which actually provides the value
- FUNCTION FMAPSH – performs, for the NC branch, the mapping in the invariant mass squared s_2 of the fermion pair defined by the flag IFERM2
- FUNCTION FMAPSZ – performs, for the NC branch, the mapping in the invariant mass squared s_1 of the fermion pair defined by the flag IFERM1
- FUNCTION FMBW – maps Breit-Wigner resonance peaks
- FUNCTION FMQED – maps peaks of the type $1/s$
- FUNCTION FOTF – computes the leptonic vacuum polarization for ALRUN from lepton and effective quark masses
- FUNCTION FOTF1 – computes the leptonic vacuum polarization for ALRUN from lepton masses and the hadronic vacuum polarization of reference [21]
- FUNCTION FSZ – performs, for the NC branch, the integration over s_2 ; this is the complement to the function FINE12
- FUNCTION FTBINV – represents the inverse function of FXTB
- FUNCTION FXSECB – provides, in dependence on the flag IDCS and for the CC03 process, the threefold differential cross-section $d^3\sigma/(dx_1 dx_2 d\cos\theta)$ or the argument $T_n(\cos\theta_{lab}) d^3\sigma_{CC03}/(ds_1 ds_2 d\cos\theta)$ of the moment integral (2.15)
- FUNCTION FXSECD – provides, in dependence on the flags IDCS and IBIN and for the CC03 process, the integral over the threefold differential cross-section $d^3\sigma/(dx_1 dx_2 d\cos\theta)$ for given limits
- FUNCTION FXTB – is the mapping function for the $\cos\theta$ integration with the subroutine FDSIMV
- FUNCTION g222 (g322, g322m, g422, g233) – computes the kinematic function $\mathcal{G}_{222}$ ($\mathcal{G}_{322}$, $\mathcal{G}_{322}^a$, $\mathcal{G}_{422}$, $\mathcal{G}_{233}$) introduced in references [8,9]
- SUBROUTINE GFBGD2 – computes the kinematic functions $\mathcal{G}_k$ for the pure background for the CC11 process
- SUBROUTINE GFBGD3 – computes the kinematic functions $\mathcal{G}_k$ for the signal-background interferences of the CC11 process
- SUBROUTINE GFBRN – computes the kinematic functions $\mathcal{G}_k$ for the CC03 process (compare equation (2.4), see reference [7])
- SUBROUTINE GFBRNA – computes, in dependence on the flags IDCS and IANO, the angular kinematic functions $\mathcal{G}_k^{\text{CC03}}(s; s_1, s_2, \cos\theta)$ (compare eq. (2.13), see

reference [5])

- SUBROUTINE GFBRNB – computes, in dependence on the flags IDCS and IANO, the integrated angular kinematic functions $\mathcal{G}_k^{\text{CC03}}(s; s_1, s_2, \cos\theta_{\min}, \cos\theta_{\max})$
- SUBROUTINE GFBRNC – computes the kinematic function for the NCqed branch
- MAIN – performs the ramification between the NCqed and the CC/NC branches of the code; carries out some initialization; calls WWIN00 and WWIN01 for further initialization; loops over the center of mass energies; loops over the desired values of IREGIM and IMOMN; prints results
- FUNCTION MYEXP1 (MYEXP2) – is the inverse function of MYLOG1 (MYLOG2) as physics input
- FUNCTION MYLOG1 (MYLOG2) – is the mapping function for the $s_1(s_2)$ integration used in calls to FDSIMP (FDSIMS)
- SUBROUTINE NCGENT – the central routine for the NCqed branch of the code; sets integration precisions; calls NCIN00 and NCIN01 for further initialization; loops over the center of mass energies; provides the total cross-section for the NCqed branch; prints results
- SUBROUTINE NCIN00 – initializes the flags needed for the NCqed branch of the code; initializes the fermion types and couplings and performs other kinematics independent initialization for the NCqed branch
- SUBROUTINE NCIN01 – performs, for the NCqed branch of the code, the initialization of quantities that depend on s only and provides the soft+virtual radiator
- SUBROUTINE NCIN02 – provides the phase space factor and the coupling function for the NCqed branch given in equation (4.5) of first of references [6]
- SUBROUTINE prop – provides the Z , γ , and gluon propagators for use in the subroutine dsdshsz
- SUBROUTINE PROPAG – computes the neutral gauge boson propagators used in NCIN02
- FUNCTION RHOINV – is the inverse function of RHOPR
- FUNCTION RHOPR – provides, for the CC/NC branch of the code and according to the mapping choice via IMAPPG, the mapping function for the s_1 and s_2 integrations
- FUNCTION RI3 – computes the scalar two-point function needed in FOTF and FOTF1
- SUBROUTINE SETANO – initializes the anomalous couplings defined by equations (B.3) to (B.8) in dependence on the flags IDCS and IANO
- SUBROUTINE SHORTI – computes the kinematic functions $\mathcal{G}_{\text{CC11}}^{uu,dd}$ and $\mathcal{G}_{\text{CC11}}^{u,d}$ as given in equations (3.9) and (3.10) of reference [5]
- SUBROUTINE SIMPS, SIMPT – performs the integration of an input function inside given limits and with a given precision by applying a self-adaptive Simpson algorithm
- FUNCTION SPENC – is an auxiliary function used by the complex dilogarithm function XSPENZ
- FUNCTION STRUCF – computes, according to the value of IZETTA, the structure function $D(x)$ from appendix A of reference [15]

- FUNCTION TOTXS – computes the total cross-section for the CC/NC branch and the differential cross-section for the CC03 case
- FUNCTION TRILOG – provides the trilogarithm function $\text{Li}_3(x)$ for $0 \leq x \leq 1$
- FUNCTION TRIS12 – provides the Nielsen function $S_{1,2}(x)$ for $0 \leq x \leq 1$
- SUBROUTINE WUFLAG – initializes and changes the default flags as well as physics input
- SUBROUTINE WWIN00 – performs the kinematics independent initialization of mathematical and physical constants for the CC/NC branch of the code; calls the subroutines `bbmmin` and `SETANO` for further initialization
- SUBROUTINE WWIN01 – performs, for the CC/NC branch, the initialization of quantities that depend on the center of mass energy or, equivalently, s only; performs initialization in dependence on the flags `IPROC`, `ICONVL`, and `IQEDHS`; provides the soft+virtual radiator for the CC/NC computation
- SUBROUTINE WWIN02 – performs the initialization of tree level quantities that depend on s , s_1 , and s_2 ; for the CC/NC branch
- SUBROUTINE WWIN03 – performs the initialization of radiative quantities that depend on s , s_1 , and s_2 ; for the CC/NC branch
- SUBROUTINE XSBRNC – provides the tree level cross-section for the NCqed branch of the code
- SUBROUTINE XSEC2E – is needed for the computation of ISR-corrected cross-sections, moments, and distributions; performs the s' integration for the flux function approach and the x_+ integration for the structure function approach
- SUBROUTINE XSEC3B – computes, in the flux function approach and for the CC11 family with `IBCKGR=1`, the universally ISR-corrected, threefold (with respect to s_1 , s_2 , and s') differential cross-section for background contributions by calling the subroutine `XSECBG`
- SUBROUTINE XSEC3E – is used in the flux function convolution for the CC03 process, i.e. for `IPROC=1`, `IBORNF=1`, and `ICONVL=0` only; computes, in dependence on the settings of `ITNONU` and `IQEDHS` and according to formula (2.6) and reference [10], the threefold differential CC03-expressions $[\beta_e v^{\beta_e-1} \mathcal{S}_k + \mathcal{H}_k]$
- SUBROUTINE XSECB – for the CC/NC branch; provides, in dependence on the flags `IPROC`, `IDCS`, and `IREGIM`, the twofold differential (with respect to s_1 and s_2) cross-section or moments for the chosen tree level process by calling `GFBRN` or integrating `FXSECB` over the W production angle
- SUBROUTINE XSECBG – provides, in dependence on the channel chosen via the flag `ICHNNL`, the twofold (with respect to s_1 and s_2) differential cross-section for the signal-background interference and the pure background in the CC11 process by calling the subroutines `GFBGD2` and `GFBGD3`
- SUBROUTINE XSNC2E – provides the twofold (with respect to s_1 and s_2) differential cross-section for the radiative NCqed cross-section by integrating over s'

- SUBROUTINE XSNC3E – computes, according to the flags ITNONU, the three-fold (with respect to s_1 and s_2) differential, radiative NCqed cross-section
- FUNCTION XSPENZ – provides the complex dilogarithm function $\text{Li}_2(z)$ by transforming the argument to the area of fast convergence

3.4 Program output

The output of GENTLE/4fan strongly depends on the user-chosen branch of the program or, in other words, on the flag settings. Four different output appearances are to be distinguished, namely

- (1) the output of total cross-sections and moments for the CC11 family
- (2) the output of angular distributions for the CC03 process
- (3) the output of the 4fan branch of the code
- (4) the output of the NCqed branch of the code

In the below subsections, we will provide and explain examples for each of the above four kinds of output. The below outputs will, at the same time, serve as test run output, where we assume that the user has used the flags and physical input as shown in the output. Any other pre-set input should remain unchanged for test run purposes.

3.4.1 Output of total cross-sections and moments for CC11

The output of GENTLE/4fan for the total cross-section and moments for the CC11 family of processes appears as follows.

```
***** This is GENTLE/4fan -- Version 2.0 *****
```

```
This is the CC branch of GENTLE/4fan for total cross-sections
```

```
FLAGS:IPROC ,IINPT ,IONSHL,IBORNF,IBCKGR,ICHNNL= 1 1 1 1 1 2
```

```
FLAGS:IGAMZS,IGAMW ,ITNONU,IQEDHS,ICOLMB,ICONVL= 1 0 0 1 2 0
```

```
FLAGS:IZERO ,IZETTA,IIQCD ,IDCS ,IANO ,IBIN = 0 1 1 0 0 0
```

```
FLAGS:IMAP ,IRMAX ,IRSTP ,IMMIN ,IMMAX = 1 1 1 1 4
```

```
EPS1 = 1.0000000000000000E-04
```

```
S1MIN,S1MAX= 0.0000E+00 0.2592E+05
```

```
S2MIN,S2MAX= 0.0000E+00 0.2592E+05
```

```
Energy (GeV) Cross-section (pb)
```

```
161.00 0.1336661D+00
```

```
MOMENTS
```

1	1	2	3	4
1	0.4753809D+00	0.3387719D+01	0.5607608D+02	0.1411261D+04


```

S1MIN,S1MAX= 0.0000E+00 0.3063E+05
S2MIN,S2MAX= 0.0000E+00 0.3063E+05
Energy (GeV) Cross-section (pb)
175.00 0.4952086D+00
MOMENTS
1 1 2 3 4
1 0.1126753D+01 0.8432371D+01 0.1030576D+03 0.1885898D+04
S1MIN,S1MAX= 0.0000E+00 0.3610E+05
S2MIN,S2MAX= 0.0000E+00 0.3610E+05
Energy (GeV) Cross-section (pb)
190.00 0.6080011D+00
MOMENTS
1 1 2 3 4
1 0.2151788D+01 0.2755849D+02 0.4972633D+03 0.1089393D+05
S1MIN,S1MAX= 0.0000E+00 0.4203E+05
S2MIN,S2MAX= 0.0000E+00 0.4203E+05
Energy (GeV) Cross-section (pb)
205.00 0.6355688D+00
MOMENTS
1 1 2 3 4
1 0.3208536D+01 0.5908307D+02 0.1485244D+04 0.4348380D+05

```

***** This is GENTLE/4fan -- Version 2.0 *****

```

This is the CC branch of GENTLE/4fan for total cross-sections
FLAGS:IPROC ,IINPT ,IONSHL,IBORNF,IBCKGR,ICHNNL= 1 1 1 1 0 2
FLAGS:IGAMZS,IGAMW ,ITNONU,IQEDHS,ICOLMB,ICONVL= 1 0 0 1 2 0
FLAGS:IZERO ,IZETTA,IIQCD ,IDCS ,IANO ,IBIN = 0 1 1 0 0 0
FLAGS:IMAP ,IRMAX ,IRSTP ,IMMIN ,IMMAX = 1 3 3 1 4
EPS1 = 1.0000000000000000E-04
S1MIN,S1MAX= 0.0000E+00 0.2592E+05
S2MIN,S2MAX= 0.0000E+00 0.2592E+05
Energy (GeV) Cross-section (pb)
161.00 0.1332829D+00
MOMENTS
3 1 2 3 4
3 -0.3052332D+01 0.2605644D+02 -0.3833234D+03 0.7596405D+04
S1MIN,S1MAX= 0.0000E+00 0.3063E+05
S2MIN,S2MAX= 0.0000E+00 0.3063E+05
Energy (GeV) Cross-section (pb)
175.00 0.4946375D+00
MOMENTS
3 1 2 3 4
3 -0.5666559D+00 0.8922053D+01 -0.9553119D+02 0.1918868D+04

```

```

S1MIN,S1MAX= 0.0000E+00 0.3610E+05
S2MIN,S2MAX= 0.0000E+00 0.3610E+05
Energy (GeV) Cross-section (pb)
190.00 0.6074997D+00
MOMENTS
3 1 2 3 4
3 -0.1004648D+00 0.9783632D+01 -0.4987223D+02 0.1646710D+04
S1MIN,S1MAX= 0.0000E+00 0.4203E+05
S2MIN,S2MAX= 0.0000E+00 0.4203E+05
Energy (GeV) Cross-section (pb)
205.00 0.6352834D+00
MOMENTS
3 1 2 3 4
3 0.1802810D+00 0.1232392D+02 0.6206403D+01 0.2242722D+04

```

First, the program identifies itself and the branch of the code for which output is produced. Then, all relevant flags are printed, before the relative precision EPS1 of the last computation is given. Finally, physics output is generated. For each center of mass energy the energy and the corresponding total cross-section are printed. Then, in the first column, integers k indicate the value of IREGIM and thus determine the physics quantity for which moments are output. The first line in the moments' output represents the degree n of the moments in the column below. In the second line, the moment of degree n for the quantity determined by k is given. Its value is normalized to the total cross-section.

3.4.2 Output of differential cross-sections for CC03

First we give an example of output for IBIN=0, with differential cross-sections computed at fixed values for the scattering angle.

```

***** This is GENTLE/4fan -- Version 2.0 *****

This is the CC branch of GENTLE/4fan for diff. cross-sections
XG= 0.10000D+00 XZ= -0.10000D+00
YG= 0.00000D+00 YZ= 0.00000D+00
ZZ= 0.00000D+00 DZ= 0.50000D-01
FLAGS:IPROC ,IINPT ,IONSHL,IBORNF,IBCKGR,ICHNNL= 1 1 1 0 0 2
FLAGS:IGAMZS,IGAMW ,ITNONU,IQEDHS,ICOLMB,ICONVL= 1 0 0 1 2 1
FLAGS:IZERO ,IZETTA,IIQCD ,IDCS ,IANO ,IBIN = 0 1 1 1 8 0
FLAGS:IMAP ,IRMAX ,IRSTP ,IMMIN ,IMMAX = 1 3 5 1 1
EPS1 = 1.0000000000000000E-04

```

S1MIN,S1MAX= 0.0000E+00 0.2592E+05
 S2MIN,S2MAX= 0.0000E+00 0.2592E+05
 Energy (GeV)= 161.00000000000000
 COSW Cross-section (pb)
 -1.000 0.0490713
 -0.800 0.0536632
 -0.400 0.0646671
 0.000 0.0794040
 0.400 0.1009377
 0.800 0.1381398
 1.000 0.1760354
 S1MIN,S1MAX= 0.0000E+00 0.3063E+05
 S2MIN,S2MAX= 0.0000E+00 0.3063E+05
 Energy (GeV)= 175.00000000000000
 COSW Cross-section (pb)
 -1.000 0.0951087
 -0.800 0.1123347
 -0.400 0.1564827
 0.000 0.2238638
 0.400 0.3411499
 0.800 0.5873924
 1.000 0.8353949
 S1MIN,S1MAX= 0.0000E+00 0.3610E+05
 S2MIN,S2MAX= 0.0000E+00 0.3610E+05
 Energy (GeV)= 190.00000000000000
 COSW Cross-section (pb)
 -1.000 0.0710393
 -0.800 0.0891843
 -0.400 0.1359378
 0.000 0.2124469
 0.400 0.3669140
 0.800 0.7949695
 1.000 1.3546129
 S1MIN,S1MAX= 0.0000E+00 0.4203E+05
 S2MIN,S2MAX= 0.0000E+00 0.4203E+05
 Energy (GeV)= 205.00000000000000
 COSW Cross-section (pb)
 -1.000 0.0512161
 -0.800 0.0678664
 -0.400 0.1099483
 0.000 0.1806793
 0.400 0.3368272
 0.800 0.8655771
 1.000 1.7575708

After the identification of the active GENTLE/4fan branch, the anomalous couplings (compare page 15 and appendix B) used in the CC03 subprocess are output, before the used flags and the relative precision of the run are printed. Next, below the print of each center of mass energy, the differential cross-sections for IBINNU fixed values of the scattering angle COSW are written to the output.

Below, we present output for IBIN=1, i.e the given differential cross-section values are integrated over bins in the scattering angle. The appearance of the output is identical to the above output for IBIN=0 except that, for each energy, the total cross-section is given right below the prints of scattering angle values and the corresponding cross-sections.

***** This is GENTLE/4fan -- Version 2.0 *****

This is the CC branch of GENTLE/4fan for diff. cross-sections

```

XG=    0.10000D+00    XZ=    0.65543D-01
YG=    0.00000D+00    YZ=    0.00000D+00
ZZ=    0.00000D+00    DZ=    0.11957D+00
FLAGS:IPROC ,IINPT ,IONSHL,IBORNF,IBCKGR,ICHNNL= 1 1 1 0 0 2
FLAGS:IGAMZS,IGAMW ,ITNONU,IQEDHS,ICOLMB,ICONVL= 1 0 0 1 2 1
FLAGS:IZERO ,IZETTA,IIQCD ,IDCS ,IANO ,IBIN  = 0 1 1 1 7 1
FLAGS:IMAP ,IRMAX ,IRSTP ,IMMIN ,IMMAX      = 1 3 5 1 1
EPS1 = 1.0000000000000000E-04
S1MIN,S1MAX= 0.0000E+00 0.2592E+05
S2MIN,S2MAX= 0.0000E+00 0.2592E+05
Energy (GeV)= 161.00000000000000
      COSW      Cross-section (pb)
-1.000 - -0.600    0.020923
-0.600 - -0.200    0.025547
-0.200 -  0.200    0.031763
 0.200 -  0.600    0.040919
 0.600 -  1.000    0.057338
      TOTAL XSEC =    0.176489
S1MIN,S1MAX= 0.0000E+00 0.3063E+05
S2MIN,S2MAX= 0.0000E+00 0.3063E+05
Energy (GeV)= 175.00000000000000
      COSW      Cross-section (pb)
-1.000 - -0.600    0.042294
-0.600 - -0.200    0.060777
-0.200 -  0.200    0.089224
 0.200 -  0.600    0.139347
 0.600 -  1.000    0.247555
      TOTAL XSEC =    0.579198

```

```

S1MIN,S1MAX=  0.0000E+00  0.3610E+05
S2MIN,S2MAX=  0.0000E+00  0.3610E+05
Energy (GeV)=  190.00000000000000
      COSW      Cross-section (pb)
-1.000 - -0.600    0.032453
-0.600 - -0.200    0.051912
-0.200 -  0.200    0.084241
 0.200 -  0.600    0.150952
 0.600 -  1.000    0.346228
      TOTAL XSEC =    0.665786
S1MIN,S1MAX=  0.0000E+00  0.4203E+05
S2MIN,S2MAX=  0.0000E+00  0.4203E+05
Energy (GeV)=  205.00000000000000
      COSW      Cross-section (pb)
-1.000 - -0.600    0.023983
-0.600 - -0.200    0.041371
-0.200 -  0.200    0.071246
 0.200 -  0.600    0.139243
 0.600 -  1.000    0.393478
      TOTAL XSEC =    0.669320

```

3.4.3 Output for the 4fan branch

***** This is GENTLE/4fan -- Version 2.0 *****

This is the 4fan branch of GENTLE/4fan
4fan was called for fermions 13 5
with masses 0.500000D-03 0.100000D-01
total cross section is calculated

```

FLAGS:IPROC ,IINPT ,IONSHL,IBORNF,IBCKGR,ICHNNL= 2 0 1 0 0 0
FLAGS:IGAMZS,IGAMW ,ITNONU,IQEDHS,ICOLMB,ICONVL= 1 0 0 1 0 0
FLAGS:IZERO ,IZETTA,IIQCD ,IDCS ,IANO ,IBIN  = 1 1 0 0 0 0
FLAGS:IMAP ,IRMAX ,IRSTP ,IMMIN ,IMMAX      = 1 0 1 1 1
FLAGS FOR 4FAN: IFERM1, IFERM2, IHIGGS=13 5 0
EPS1 = 1.0000000000000000E-04
ABS1 = 1.0000000000000000E-04
S1MIN,S1MAX=  0.5805E+04  0.1128E+05
S2MIN,S2MAX=  0.9000E+03  0.2592E+05

```

Energy (GeV)	Cross-section (pb)
161.00	0.2907564D+00
S1MIN,S1MAX=	0.5805E+04 0.1128E+05
S2MIN,S2MAX=	0.9000E+03 0.3063E+05
Energy (GeV)	Cross-section (pb)
175.00	0.8156756D+00
S1MIN,S1MAX=	0.5805E+04 0.1128E+05
S2MIN,S2MAX=	0.9000E+03 0.3610E+05
Energy (GeV)	Cross-section (pb)
190.00	0.9510082D+01
S1MIN,S1MAX=	0.5805E+04 0.1128E+05
S2MIN,S2MAX=	0.9000E+03 0.4203E+05
Energy (GeV)	Cross-section (pb)
205.00	0.1249948D+02

3.4.4 Output for the NCqed branch

***** This is GENTLE/4fan -- Version 2.0 *****

This is the NCqed branch of GENTLE/4fan
 FLAGS: NINPT,NNCPRC,NGAMZS,NBORNF,NTNONU,NQEDHS
 0 2 1 1 0 0

GFER = 1.1663900000000000E-05
 AMW = 80.23
 AMZ = 91.1888
 GAMZ = 2.4974
 ALPW = 7.808229874287499E-03
 SIN^2(\theta_W) = .2310309124515784

IFERM1,IFERM2 = 2 4
 RNCOU1,RNCOU2 = 1.00000000 3.00000000
 AM1 ,AM2 = .105658389 4.3
 CUTM12,CUTM34 = .211316778 8.6
 CUTXPR = .0

EPS1 1.0000000000000000E-05
 ABS1 1.0000000000000000E-15
 ENERGY (GeV) CROSS-SECTION (fb)
 161.0 .8105861091D+02
 175.0 .6725488126D+02
 190.0 .6303702520D+02
 205.0 .5842607173D+02

Table 3

Correspondence between fermion pair indices and fermion types for the NCqed branch of GENTLE/4fan.

Fermion Pair Index	1	2	3	4
Fermion Type	ν_μ, ν_τ	μ, τ	u, c	d, s, b

After identifying the active branch, the flags relevant to the NCqed branch of the code are printed. The flags are as set in WUFLAG, but the first letter is changed to an N standing for NCqed. Next, the relevant physical input constants are printed the meaning of which is described in section 3.2. The next paragraph of output begins with the fermion pair indices. The correspondence of fermions and indices is given in table 3. Then, for the two fermion types used in the computation, the color factors (RNCOU1, RNCOU2), the fermion masses (AM1, AM2), and the lower invariant mass cuts (CUTM12, CUTM34) are printed. Closing this paragraph of output, the minimum value CUTXPR of the fraction s'/s is printed. Finally, together with the relative and absolute required precisions EPS1 and ABS1, the total cross-sections are output for each required center of mass energy.

3.5 Use of the program

In this section, we will give a short guide to the compilation and use of GENTLE/4fan.

We recommend to exhaust all means of compiler optimization, because, according to our experience, the CPU time consumption can thus be reduced by a factor of two or more. When available, the use of a compiler option that makes the default size of floating-point constants REAL*8 may be advantageous. For HP computers, we recommend

```
f77 +04 +Onolimit +ppu -K -R8 gentle.f .
```

For Silicon Graphics computers, compilation with

```
f77 -r8 -03 -non_shared -G 32 -jmpopt -mips2 -static gentle.f
```

has proved successful.

Unless the user wants to run GENTLE/4fan with default flags, the user should set the flags described in section 3.2. The setting of a flag IFLAG to a value IVALUE is accomplished by adding the line

```
CALL WUFLAG('IFLAG', IVALUE)
```

after the `DATA` statements in the `MAIN` of `GENTLE/4fan`. Please note that the numerical format of `IVALUE` is `INTEGER`. Similarly, for `IANO=8`, the anomalous couplings `XG`, `YG`, `XZ`, `YZ`, `ZZ`, `DG`, and `DZ` may be set by calls like

```
CALL WUFLAG('XG', VALUE)
```

at the same place in the `MAIN`. Please note that the numerical format of `VALUE` is `DOUBLE PRECISION`. In addition, the physical input variables `GFER`, `ALPW`, `AME`, `AMW`, `AMZ`, `GAMZ`, and `ALPHS` for the `CC` branch of the code as well as `AMHIG` and `GAMHIG` for the `4fan` branch can be changed by analogous calls to `WUFLAG`. Finally, it is possible to change the physics input to the `4fan` and `NCqed` branches of the code in the subroutines `bbmmin` and `NCIN00` respectively.

To determine the physics quantities and the degrees of moments to be computed and printed, the user must set the range of the loops over `IREGIM` and `IMOMN` (see page 18).

Cuts on the final state invariant pair masses may be applied in the `CC` and the `4fan` branches of the code by altering `S1MIN`, `S1MAX`, `S2MIN`, and `S2MAX` in the `MAIN`. In the `NCqed` branch, final state invariant pair mass cuts are implemented by setting `X12MIN`, `X12MAX`, `X34MIN`, and `X34MAX` in the subroutines `NCGENT` and `FINC34` ($X12 = s_1/s$, $X34 = s_2/s$).

Acknowledgments

We gratefully acknowledge the valuable contributions of M. Bilenky in early phases of the project. We benefitted greatly from discussions with and extensive tests of the program by Peter Clarke, Douglas Ferguson, Som Ganguli, Bingnian Jin, Martin Grünewald. We would also like to thank A. Ballestrero, F.A. Berends, M. Dubinin, S. Jadach, M. Moretti, O. Nicrosini, T. Ohl, G. Passarino, R. Pittau, M. Pohl, M. Skrzypek, and Z. Was for kind and productive collaboration during the LEP 2 and LC workshops in 1995 and 1996.

A Numerical Integration in GENTLE/4fan

The integrations over s_1, s_2 are performed numerically by a successive use of an adaptive one-dimensional Simpson integration routine. One (two) more integrations are needed to calculate initial state QED corrections in the flux function (structure function) approach. Mappings of the integrands may be chosen with flag **IMAP**. We now describe the mappings.

After the analytical integration over the six angular variables, one has to treat Breit-Wigner resonances arising from the exchange of intermediate bosons:

$$f_{BW}(V, s) \sim \frac{1}{(s - M_V^2)^2 + \Gamma_V^2 M_V^2}, \quad V = Z, W, H. \quad (\text{A.1})$$

In the case of neutral current processes, additional peaks appear at small momentum transfer due to virtual photons or gluons,

$$f_{low}(V, s) \sim \frac{1}{s}, \quad V = \gamma, g. \quad (\text{A.2})$$

For charged current processes, the standard mapping of the Breit-Wigner resonance of the W boson is applied in **GENTLE/4fan**.

In the case of neutral current processes, the peaks at small momentum transfer and the Breit-Wigner resonances of the Z and the Higgs boson have to be mapped simultaneously. In **GENTLE/4fan**, the singularity at small momentum transfer is mapped first by the usual transformation to a new variable $\bar{s}$,

$$\bar{s} = f_1(s) = \ln s. \quad (\text{A.3})$$

Then follows a mapping of the Z resonance, which does not destroy the previous mapping,

$$\bar{\bar{s}} = f_2(\bar{s}) = \bar{s} + \frac{c}{[\bar{\Gamma}_V \bar{M}_V]} \arctan \frac{\bar{s} - \bar{M}_V^2}{[\bar{\Gamma}_V \bar{M}_V]}. \quad (\text{A.4})$$

The variables $\bar{M}_V^2$ and $[\bar{\Gamma}_V \bar{M}_V]$ are calculated taking into account the previous mapping,

$$\bar{M}_V^2 = f_1(M_V^2), \quad [\bar{\Gamma}_V \bar{M}_V] = f_1(M_V^2 + \Gamma_V M_V) - f_1(M_V^2). \quad (\text{A.5})$$

The function f_2 is inverted numerically by Newton's Method,

$$\bar{\bar{s}}_{i+1} = \bar{\bar{s}}_i - \frac{f_2(\bar{\bar{s}}_i)}{f_2'(\bar{\bar{s}}_i)}, \quad (\text{A.6})$$

which always converges for the starting value $\bar{\bar{s}}_0 = \bar{M}_V^2$. The free constant c

in equation (A.4) is optimized for every final state separately. However, the quality of the mapping (A.4) is sensitive to the order of magnitude of c only. In the case of Higgs production, a procedure similar to (A.4) is added to map the Higgs resonance.

B Anomalous Triple-Boson Couplings

In addition to the SM Lagrangian, we consider the following CP conserving operators of dimension 6¹⁰:

$$\begin{aligned} \Delta\mathcal{L}_{\text{eff}} = & g' \frac{\alpha_{B\phi}}{m_W^2} (D_\mu \Phi)^\dagger B^{\mu\nu} (D_\nu \Phi) + g \frac{\alpha_{W\phi}}{m_W^2} (D_\nu \Phi)^\dagger \overleftrightarrow{\tau} \overleftrightarrow{W}^{\mu\nu} (D_\nu \Phi) \\ & + g \frac{\alpha_W}{6m_W^2} \overleftrightarrow{W}_\nu^\mu (\overleftrightarrow{W}_\rho^\nu \times \overleftrightarrow{W}_\mu^\rho). \end{aligned} \quad (\text{B.1})$$

In the unitary gauge, these operators lead to the following effective Lagrangian for the WWV vertex:

$$\begin{aligned} \mathcal{L}_{\text{eff}}^{WWV} = & ig_{WWV} \left[g_1^V (W_{\mu\nu}^+ W^{-\mu} - W^{+\mu} W_{\mu\nu}^-) V^\nu \right. \\ & \left. + \kappa_V W_\mu^+ W_\nu^- V^{\mu\nu} + \frac{\lambda_V}{m_W^2} W_\mu^{+\nu} W_\nu^{-\rho} V_\rho^\mu \right], \end{aligned} \quad (\text{B.2})$$

where $V \in \{Z, \gamma\}$. Therefore, one has six relevant anomalous triple-boson couplings, namely g_1^γ , g_1^Z , κ_γ , κ_Z , λ_γ , and λ_Z . Electromagnetic gauge invariance requires $g_1^\gamma = 1$. In the HISZ-scenario [20] ($\text{I\!A\!N\!O} = \pm 7$), one sets $\alpha_{B\phi} = \alpha_{W\phi}$. Thus, in this scenario, one obtains relations between the parameters κ_γ , κ_Z , and g_1^Z .

For the WWZ -vertex, we also include a C and P violating term in the Lagrangian:

$$\mathcal{L}_Z = - \frac{eZ_Z}{m_W^2} \partial_\alpha \hat{Z}_{\rho\sigma} \left(W^{+\alpha} \overleftrightarrow{\partial}^\rho W^{-\sigma} - W^{+\sigma} \overleftrightarrow{\partial}^\rho W^{-\alpha} \right),$$

where we have introduced the dual field strength tensor

$$\hat{Z}_{\rho\sigma} = \frac{1}{2} \epsilon_{\rho\sigma\alpha\beta} Z^{\alpha\beta}.$$

To disentangle the anomalous couplings' contributions to the cross-section, we

¹⁰ For definitions of anomalous couplings, see e.g. [22,16].

use redefined parameters:

$$\delta_Z = (g_1^Z - 1) \cot \theta_W, \quad (\text{B.3})$$

$$x_\gamma = \kappa_\gamma - 1, \quad (\text{B.4})$$

$$x_Z = (\kappa_Z - 1)(\cot \theta_W + \delta_Z), \quad (\text{B.5})$$

$$y_\gamma = \lambda_\gamma, \quad (\text{B.6})$$

$$y_Z = \lambda_Z \cot \theta_W, \quad (\text{B.7})$$

$$z_Z = z_Z. \quad (\text{B.8})$$

The **GENTLE/4fan** user can set values for the anomalous couplings defined in equations (B.3) to (B.8). In the **SM**, the anomalous couplings (B.3) to (B.8) vanish. The special case of anomalous couplings induced by a heavy extra neutral gauge boson is described in appendix C.

C Treatment of Extra Neutral Gauge Bosons

The effect of an extra neutral gauge boson Z' can be described with **GENTLE/-4fan** by two anomalous couplings $g_{WW\gamma}^*$ and g_{WWZ}^* of the photon and the Standard Model Z boson to W pairs [23],

$$g_{WW\gamma}^* = 1 + \delta_\gamma \quad \text{and} \quad g_{WWZ}^* = \cot \theta_W + \delta_Z, \quad (\text{C.1})$$

where

$$\begin{aligned} \delta_\gamma &= g_{WWZ_1} \left(\frac{a_1}{a} - \frac{v_1}{v} \right) v(1 + \Delta\chi)\chi_Z + g_{WWZ_2} \left(\frac{a_2}{a} - \frac{v_2}{v} \right) v\chi_2, \\ \delta_Z &= -\cot \theta_W + g_{WWZ_1} \frac{a_1}{a} (1 + \Delta\chi) + g_{WWZ_2} \frac{a_2}{a} \frac{\chi_2}{\chi_Z}, \end{aligned} \quad (\text{C.2})$$

and

$$\chi_Z = \frac{s}{s - M_Z^2}, \quad \Delta\chi = -\frac{2M_Z(M_Z - M_1)}{s - M_Z^2}, \quad \chi_2 = \frac{s}{s - M_2^2}. \quad (\text{C.3})$$

In equations (C.2) and (C.3), M_i, v_i, a_i , $i = 1, 2$ denote the masses and electron couplings of the mass eigenstates Z_1 and Z_2 , which are, in general, a result of a mixing of the gauge symmetry eigenstates Z and Z' ,

$$\begin{aligned}
Z_1 &= Z \cos \theta_M + Z' \sin \theta_M, & Z_2 &= -Z \sin \theta_M + Z' \cos \theta_M, \\
v_1 &= v \cos \theta_M + v' \sin \theta_M, & v_2 &= -v \sin \theta_M + v' \cos \theta_M, \\
a_1 &= a \cos \theta_M + a' \sin \theta_M, & a_2 &= -a \sin \theta_M + a' \cos \theta_M,
\end{aligned} \tag{C.4}$$

where v, a (v', a') denote the couplings of the symmetry eigenstates Z (Z') to electrons and θ_M is the ZZ' mixing angle. The couplings g_{WWZ_1} and g_{WWZ_2} are fixed by the condition that only the Z couples to W pairs,

$$g_{WWZ_1} = \cot \theta_W \cos \theta_M, \quad g_{WWZ_2} = \cot \theta_W \sin \theta_M. \tag{C.5}$$

The mass shift from M_Z to M_1 in the Z_1 propagator is absorbed into δ_γ and δ_Z through $\Delta\chi$ assuming $M_Z - M_1 \ll M_Z$ and $M_Z - M_1 \ll (s - M_Z^2)/(2M_Z)$. Both approximations are equivalent for W pair production.

D Options of GENTLE

Table D.1 shows the observables GENTLE can calculate and the options, which may be chosen. Please note that in the CC11 family anomalous couplings are included only for the CC03 process.

Table D.1

Options of GENTLE; SF - structure function approach; FF - flux function approach; h.o. - higher order QED corrections; Coul. - Coulomb singularity.

	IREGIM	on-shell	Born	SF	FF	h.o.	Coul.	non-uni.	anom.c.
σ^{CC03}	0	+	+	+	+	+	+	+	+
σ^{CC03}	1	+	−	+	+	+	+	+	+
σ^{CC03}	2	+	−	+	−	+	+	−	+
σ^{CC03}	3	+	−	+	+	+	+	+	+
σ^{CC03}	4	+	+	+	−	+	+	−	−
$\frac{d\sigma^{\text{CC03}}}{d\cos\theta}$	−	+	+	+	−	+	+	−	+
σ^{CC11}	0	−	+	+	+	+	+	−	+
σ^{CC11}	1	−	−	+	+	+	+	−	+
σ^{CC11}	2	−	−	+	−	+	+	−	+
σ^{CC11}	3	−	−	+	+	+	+	−	+
σ^{NC02}	−	−	+	−	+	+	−	+	−
σ^{NC08}	−	−	+	−	+	+	−	+	−
σ^{NC32}	−	−	+	+	+	+	−	−	−

References

- [1] D. Bardin, R. Kleiss et al., *Report of the Working Group on Event Generators for WW Physics*, in G. Altarelli, T. Sjöstrand, F. Zwirner, eds., *Proceedings of the Workshop on Physics at LEP 2*, CERN Yellow Report CERN 96-01 (1996), Vol. 2, 3-101, and references therein.
- [2] F. Boudjema, B. Mele et al., *Report of the Working Group on Standard Model Processes*, in G. Altarelli, T. Sjöstrand, F. Zwirner, eds., *Proceedings of the Workshop on Physics at LEP 2*, CERN Yellow Report CERN 96-01 (1996), Vol. 1, 207-248, and references therein.
- [3] D. Bardin, M. Bilenky, D. Lehner, A. Olchevski and T. Riemann, *Semi-analytical approach to four-fermion production in e^+e^- annihilation*, in: T. Riemann and J. Blümlein (eds.), *Proc. of the Zeuthen Workshop on Elementary Particle Theory "Physics at LEP200 and Beyond"*, *Nucl. Phys. B* (Proc. Suppl.) **37B** (1994), 148-157.
- [4] F.A. Berends, R. Pittau and R. Kleiss, *Nucl. Phys.* **B424** (1994) 308.
- [5] D. Bardin and T. Riemann, *Nucl. Phys.* **B462** (1996) 3.
- [6] D. Lehner, Ph.D. thesis, Humboldt-Universität zu Berlin, Germany (1995), Internal Report DESY-IfH 95-07, e-Print Archive: hep-ph/9512301;
D. Bardin, D. Lehner and T. Riemann, *Nucl. Phys.* **B477** (1996) 27.
- [7] T. Muta, R. Najima and S. Wakaizumi, *Mod. Phys. Lett.* **A1** (1986) 203.
- [8] D. Bardin, A. Leike and T. Riemann, *Phys. Lett.* **B353** (1995) 513.
- [9] D. Bardin, A. Leike and T. Riemann, *Phys. Lett.* **B344** (1995) 383.
- [10] D. Bardin, M. Bilenky, A. Olchevski and T. Riemann, *Phys. Lett. B* **308**, (1993) 403; E: *ibid*, *B* **357**, (1995) 725.
- [11] D. Bardin, W. Beenakker and A. Denner, *Phys. Lett.* **B317** (1993) 213.
- [12] V.S. Fadin, V.A. Khoze, A.D. Martin and A. Stirling, *Phys. Lett.* **B363** (1995) 112.
- [13] Particle Data Group, R.M. Barnett et al., *"Review of Particle Physics"*, *Phys. Rev.* **D54** (1996).
- [14] V.S. Fadin, V.A. Khoze and A.D. Martin, *Phys. Lett.* **B311** (1993) 311.
- [15] F.A. Berends, W. Beenakker et al., *Report of the Working Group on WW cross-sections and distributions*, in G. Altarelli, T. Sjöstrand, F. Zwirner, eds., *Proceedings of the Workshop on Physics at LEP 2*, CERN Yellow Report CERN 96-01 (1996), Vol. 1, 79-139, and references therein.
- [16] G. Gounaris, J.-L. Kneur and D. Zeppenfeld et al., *Report of the Working Group on Triple Gauge Boson Couplings*, in G. Altarelli, T. Sjöstrand, F. Zwirner, eds., *Proceedings of the Workshop on Physics at LEP 2*, CERN Yellow Report CERN 96-01 (1996), Vol. 1, 525-576, and references therein.

- [17] J. Biebel and T. Riemann, in preparation.
- [18] D. Bardin, M. Bilenky, A. Olchevski and T. Riemann, FORTRAN code `GENTLE_1993`, unpublished;
D. Bardin, M. Bilenky, D. Lehner, A. Leike, A. Olchevski and T. Riemann, FORTRAN code `GENTLE/4fan_1995`; see also [1], p.26, and [24], p.304, and D. Bardin, D. Lehner, A. Leike and T. Riemann, Internal Report DESY-Zeuthen 96-05 (1996) [hep-ph/9603438].
- [19] A. Leike, preprint LMU-06-96, to appear in: T. Riemann and J. Blümlein, eds., *Proc. of the Zeuthen Workshop on Elementary Particle Theory “QED and QCD in higher orders”*, Rheinsberg, Germany, April 1996, e-Print Archive: hep-ph/9606262.
- [20] K. Hagiwara, S. Ishihara, R. Szalapski and D. Zeppenfeld, *Phys. Lett.* **B283** (1992) 353 and *Phys. Rev.* **D48** (1993) 2182.
- [21] S. Eidelman and F. Jegerlehner, *Z. Phys.* **C67** (1995) 585.
- [22] K. Hagiwara, K. Hikasa, R. Peccei and D. Zeppenfeld, *Nucl. Phys.* **B282** (1987) 253.
- [23] A. Pankov and N. Paver, ICTP Trieste preprint IC/96/175 [hep-ph/9610509].
- [24] M. Mangano, R. Ridolfi et al., *Report of the Working Group on Event Generators for Discovery Physics*, in G. Altarelli, T. Sjöstrand, F. Zwirner, eds., *Proceedings of the Workshop on Physics at LEP 2*, CERN Yellow Report CERN 96-01 (1996), Vol. 2, 299-353.